



OPEN ACCESS

EDITED BY

Dezhi Wu,
Hunan Agricultural University, China

REVIEWED BY

Bohan Liu,
Hunan Agricultural University, China
Peng Zhang,
China National Rice Research Institute
(CAAS), China
Huaiyong Luo,
Oil Crops Research Institute (CAAS),
China

*CORRESPONDENCE

Feng Wang,
fwang1631@163.com

SPECIALTY SECTION

This article was submitted to Plant
Genomics,
a section of the journal
Frontiers in Genetics

RECEIVED 08 September 2022

ACCEPTED 26 October 2022

PUBLISHED 11 November 2022

CITATION

Huo X, Xiao J, Peng X, Lin Y, Liu D, Liu W,
Liao Y, Li J, Zhu M, Fu C, Zeng X, Ma X,
Kong L and Wang F (2022), The grain
yield regulator *NOG1* plays a dual role in
latitudinal adaptation and cold tolerance
during rice domestication.
Front. Genet. 13:1039677.
doi: 10.3389/fgene.2022.1039677

COPYRIGHT

© 2022 Huo, Xiao, Peng, Lin, Liu, Liu,
Liao, Li, Zhu, Fu, Zeng, Ma, Kong and
Wang. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License](#)
(CC BY). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

The grain yield regulator *NOG1* plays a dual role in latitudinal adaptation and cold tolerance during rice domestication

Xing Huo¹, Junyi Xiao^{1,2}, Xin Peng¹, Yanhui Lin³, Dilin Liu¹,
Wuge Liu¹, Yilong Liao¹, Jinhua Li¹, Manshan Zhu¹,
Chongyun Fu¹, Xueqin Zeng¹, Xiaozhi Ma¹, Le Kong¹ and
Feng Wang^{1*}

¹Guangdong Key Laboratory of New Technology in Rice Breeding, Guangdong Rice Engineering Laboratory, Rice Research Institute, Guangdong Academy of Agricultural Sciences, Guangzhou, China, ²College of Agriculture, South China Agricultural University, Guangzhou, China, ³Hainan Scientific Research Station of Crop Gene Resource and Germplasm Enhancement, Ministry of Agriculture, Hainan Key Laboratory of Crop Genetics and Breeding, Institute of Food Crops, Hainan Academy of Agricultural Sciences, Haikou, China

Rice originated in tropical and subtropical regions and is distributed worldwide. Low temperature is one of the most critical abiotic stresses affecting grain yield and geographical distribution of rice. It is vital to elucidate the molecular mechanism of chilling tolerance in rice for ensuring cereals production. Previously we isolated the domestication-related gene *NOG1* which affects rice grain number and yield. In this study, we specified that rice varieties harboring high-yielding *NOG1* allele are more distributed in low-latitude regions. Additionally, we observed *NOG1* influences the chilling tolerance of rice. Through genome-wide transcriptional analysis after cold treatment at 10°C, there were 717 differentially expressed genes (DEGs) in *nog1* near-isogenic lines compared with the control Guichao 2, including 432 up-regulated DEGs and 284 down-regulated DEGs. Gene ontology annotations and KEGG enrichment analysis of DEGs showed that various biological processes and signaling pathways were related to cold stress, such as lipid metabolism and genetic information processing. These results provide new insights into the mechanism of chilling tolerance in rice and the molecular basis of environmental adaptation during rice domestication.

KEYWORDS

NOG1, cold stress, domestication, differentially expressed genes, RNA-seq

Introduction

Rice is one of the most important food crops, feeding more than half of the world's population (Ito and Lacerda, 2019). Coping with various biotic and abiotic stresses and maintaining stable yields during rice production is of great significance to ensuring world food security (Lesk et al., 2016). Low temperature is an essential abiotic stress factor




FIGURE 1
Geographical distribution of rice varieties containing the diverse *NOG1* alleles.

affecting the environmental adaptability and geographical distribution of rice. Since rice originated in tropical and subtropical regions, it is extremely sensitive to low temperature. The damage caused by low temperature will reduce the growth and development of rice, and eventually lead to the reduction of yield and quality (Jaglo et al., 2001; Zhang et al., 2019). Cold stress is prevalent in the world's main rice producing areas, and leads to huge economic losses every year. There are more than 15 million hectares of rice in the world facing the threat of low temperature (Lin et al., 2004; Zhang et al., 2014; Pradhan et al., 2019; Li et al., 2021). As an essential abiotic stress, cold stress has been a primary spot in plant science research. A series of cold tolerance-related genes were described (Saito et al., 2004; Fujino et al., 2008; Fujino and Matsuda, 2010; Su et al., 2010; Liu et al., 2013; Ma et al., 2015; Manishankar and Kudla, 2015; Shi and Gong, 2015; Mao et al., 2019). However, the molecular mechanisms of cold stress tolerance in rice are still unclear. In recent years, RNA-Seq has been widely used to perform genome-wide gene expression analysis, providing detailed and in-depth data for the molecular mechanism study of plant cold tolerance (Bashir et al., 2019; Wai et al., 2021). *NOG1* is a functional gene that regulates grain number and yield in rice. The functional site of *NOG1* is a 12-bp copy number variation in the promoter region. There are two haplotypes of *NOG1* in cultivated rice, with two copies of 12-

bp fragments in high-yielding rice and only one copy of 12-bp in low-yielding rice (Huo et al., 2017). In this study, we found that the yield-related *NOG1* is associated with the geographic distribution of rice varieties and acts as a negative player in cold tolerance during rice domestication. The genome-wide transcriptional identification of *NOG1* near-isogenic lines under cold stress were additionally compared, which provided new insights into the molecular mechanism of *NOG1*-mediated cold tolerance pathway in rice.

Results

Geographical distribution pattern of *NOG1* alleles

Geographic distribution of different *NOG1* alleles were investigated using 158 rice accessions, including 84 *indica* and 74 *japonica* varieties (Supplementary Tables S1,S2). As a result, high-yielding rice varieties containing two copies of 12-bp fragment were more frequently distributed in low-latitude regions (Figure 1). Under the control of other variables, the relationship between latitude and temperature is inversely proportional, that is, the higher the latitude, the lower the temperature. The principle is that the altitude of the Sun




FIGURE 2 Genotypes and phenotypes of the *nog1* near-isogenic line. **(A)**: Genotypes of the *nog1* near-isogenic line (NIL176), SIL176 and Guichao 2. **(B–D)**: Phenotypic comparison between SIL176 and Guichao 2. **(E–G)**: Phenotypic comparison between NIL176 and Guichao 2.

decreases with increasing latitude, resulting in a decrease in solar radiation with increasing latitude, resulting in a decrease in temperature with increasing latitude. The geographic distribution of *NOG1* alleles suggests that *NOG1* might affect the response to cold stress in addition to its role in improving rice yield.

Construction of *NOG1* near-isogenic line

In order to further eliminate the interference of genetic background, SIL176, an introgression line harboring *nog1* allele in the background of Guichao 2, was selected to

backcross with Guichao 2 and then selfed. Combined with genotype identification in the BC₂F₂ population, a near-isogenic line with only one fragment introgressed in the genomic region of *nog1* compared with Guichao 2 was screened and named NIL176. Gene sequencing validated that the *NOG1* sequence in NIL176 was identical to that in SIL176. Compared with Guichao 2, NIL176 has a 12-bp deletion and 15 SNPs in the promoter region (Figure 2A), the grain number per panicle and grain yield of the near-isogenic lines were investigated, and the results showed that NIL176 exhibited a similar performance as SIL176. Compared with Guichao 2, both NIL176 and SIL176 exhibited less grains per panicle and lower yield (Figures 2B–G). Therefore, in the further study of the




FIGURE 3
Phenotypes of chilling tolerance between NIL176 and Guichao 2. **(A)**: Phenotypic comparison of NIL176 and Guichao 2 at 28°C. **(B)**: Phenotypic comparison of NIL176 and Guichao 2 after cold stress.





FIGURE 4
Comparisons of differentially expressed genes between NIL176 and Guichao 2 under cold stress. **(A)**: Numbers of differentially expressed genes in NIL176 and Guichao 2. **(B)**: Up-regulated differentially expressed genes in NIL176 and Guichao 2. **(C)**: Down-regulated differentially expressed genes in NIL176 and Guichao 2.



function of *NOG1* gene and its response to cold stress, using near-isogenic lines as materials can eliminate the interference of genetic background and obtain more reliable results.

Altered resistance of *NOG1* near-isogenic line to cold stress

To verify whether *NOG1* affects the response of rice to cold stress, NIL176 and Guichao 2 were used for further study. Rice seedlings were treated under low temperature stress at 10°C for 10 days and recovered at 28°C for three days to examine the phenotype. After cold stress, NIL176 still maintained a normal growth, while Guichao 2 showed obvious cold injury phenotypes including wilting and yellowing (Figure 3). The results showed that compared with Guichao 2, the tolerance of NIL176 to cold stress was significantly improved.

RNA sequencing and differentially expressed genes in response to cold stress

RNA-seq analysis was performed for Guichao 2 and NIL176, which were subjected to low temperature treatment at 10°C for 12 h. The differentially expressed genes (DEGs) were screened with the difference of gene expression levels more than 2 folds or

less than 0.5 fold. After cold treatment, NIL176 had 5,954 up-regulated DEGs and 5,612 down-regulated DEGs, while Guichao 2 had 4,499 up-regulated DEGs and 5,850 down-regulated DEGs (Figure 4A). NIL176 and Guichao 2 shared 3,463 up-regulated genes and 3,953 down-regulated genes (Figures 4B,C). Under cold stress, the number of down-regulated genes in NIL176 and Guichao 2 were highly close. However, the up-regulated genes in NIL176 were significantly more than those in Guichao 2. A total of 2,491 up-regulated genes were unique in NIL176.

Functional classification of data and differentially expressed genes

After cold treatment, there are 717 DEGs when compared NIL176 with Guichao 2, of which 432 were up-regulated and 285 were down-regulated (Figure 5 and Supplementary Table S3). In order to further analyze the molecular mechanism of *NOG1*-mediated response to cold stress in rice, Gene ontology (GO) annotation and KEGG pathway enrichment analysis were conducted to identify cold stress-related pathways using DEGs between NIL176 and Guichao 2. The up-regulated DEGs were involved in biological processes such as cellular response to stress, response to stimulus, and defense response. It also affects cellular components including integral component of membrane and intrinsic component of membrane (Figure 6A). Down-regulation DEGs involves biological processes such as external condensing structure, response to abiotic stimulus, and response to inorganic substance (Figure 6B).

KEGG enrichment analysis showed that up-regulated DEGs were enriched to lipid metabolism, genetic information processing, protein families: signaling and cellular processes, protein kinases and other pathways (Figure 7A), and down-regulated DEGs were enriched to genetic information processing, carbohydrate metabolism, energy metabolism, and other pathways (Figure 7B). Notably, pathways such as genetic information processing, lipid metabolism, protein families: signaling and cellular processes can be enriched in both up-regulated and down-regulated DEGs, indicating that these pathways may play an important role in the *NOG1*-mediated rice cold tolerance gene network.

Fatty acid related gene involved in *NOG1*-mediated cold stress

Fatty acids, the major constituents of membrane glycerolipids, have important biological functions for stress response in plants. In response to a decrease in ambient temperature, plants increase the level of their unsaturated fatty acids, maintaining the appropriate fluidity of membrane lipids (Sakamoto and Murata, 2002). The related genes of fatty




FIGURE 6

GO enrichment of differentially expressed genes between NIL176 and Guichao 2. (A): GO enrichment of up-regulated genes in NIL176 compared with Guichao 2. (B): GO enrichment of down-regulated genes in NIL176 compared with Guichao 2.

acid metabolism were enriched in DGEs of Guichao 2 and NIL176 (Figure 7). To investigate whether fatty acids involved in *NOG1*-mediated cold stress, we detected expression changes among fatty acids related genes (Figure 8). In the KEGG pathway of fatty acid biosynthesis, most of genes were down-regulated after cold treatment both in Guichao 2 and NIL176. However, seven genes are relatively high expression in NIL176 compared with Guichao 2 under normal condition, which suggests that NIL176 accumulates more fatty acids before cold stress, compared with Guichao 2. Long-chain acyl-CoA synthetase (*LACS*) plays a critical role in plant development and stress responses (Zhao et al., 2021). Intriguingly, the expression of *LACS* (K01897 [EC:6.2.1.3]) was particularly high expression in the cold-treated NIL176 plant.

Discussion


NOG1 as a player in rice domestication and latitude adaptability

Rice is one of the earliest crops domesticated by humans. A series of key domestication traits were artificially selected, such as seed shattering, plant architecture, grain color and awn development (Konishi et al., 2006; Li et al., 2006; Sweeney et al., 2006; Tan et al., 2008; Zhu et al., 2011; Zhu et al., 2013;

Hua et al., 2015). At the same time, rice originated in tropical and subtropical regions. During the process of domestication, local varieties need a series of adaptive improvements to the local environment, including photoperiod, temperature tolerance, drought and other stress resistance. In this study, the domestication-related *NOG1* gene affects the cold tolerance of rice, thereby affecting the latitudinal distribution of *NOG1* alleles. In the future molecular breeding process for genetic improvement using domesticated genes such as *NOG1*, it is necessary to comprehensively consider the geographic location and temperature of the planting area.

Fatty acid metabolism pathway contributes to stress resistance in rice


Biofilm is the main damaged part of plants under cold stress, and the damage can lead to irreversible phase transition of the biofilm system. The fluidity and stability of biofilms are closely related to the low temperature tolerance of rice (Sakamoto and Murata, 2002). Fatty acids are important components of biofilms, and the content and composition of unsaturated fatty acids play an influential role in the cold tolerance of rice (Ariizumi et al., 2002). The *NOG1* gene that regulates grain number and yield in rice encodes enoyl-CoA hydratase/isomerase (ECH), a key enzyme in fatty acid metabolism, which affects the



content of endogenous fatty acids and the composition of unsaturated fatty acids in rice, especially C18:3. The fatty acid biosynthesis pathway was enriched in DEGs, which indicates that fatty acid metabolism likely played an essential role in *NOG1*-mediated low temperature stress. Lipid metabolism, alpha-Linolenic acid metabolism, fatty acid biosynthesis and fatty acid degradation pathways were all enriched.

Cold stress response pathways are involved in various biological processes

RNA-seq can analyze the differential expression of genome-wide genes, which is helpful to comprehensively analyze the response pathway of rice cold tolerance. The results of GO annotation and KEGG enrichment analysis of DEGs showed that in addition to fatty acid metabolism pathways, there are



multiple pathways involved in cold stress responses. GO annotation enriched in transmembrane transporter activity, integral component of membrane, intrinsic component of membrane and response to inorganic substance. Membrane transporters are important functional proteins that change biomembrane permeability, maintain osmotic pressure and

ion transport inside and outside the biomembrane, and are one of the essential ways to improve plant cold tolerance (Lissarre et al., 2010; Miura and Furumoto, 2013).

Additionally, cold stress is related to plant stress resistance (Saijo and Loo, 2020). In this study, GO analysis showed that stress resistance-related pathways such as external encapsulating structure, response to abiotic stimulus, and response to inorganic substance were activated. The results of KEGG also showed that environmental adaptation, environmental adaptation pathways such as Information processing and plant-pathogen interaction are enriched. Hence, *NOG1* may also be related to other abiotic stresses and disease resistance in rice in addition to its role in low temperature stress. These results enrich the understanding of cold resistance pathways in rice and provide new evidence for analyzing the molecular mechanism of rice cold tolerance.

Conclusion

The *NOG1* gene, which regulates grain number and yield in rice, encodes a key enzyme in fatty acid metabolism. Allele identification and geographic origin of 158 rice varieties including 84 indica varieties and 74 japonica varieties indicated a strong latitude adaptation of the *NOG1* alleles. Through the cold treatment of Guichao 2 and NIL176 (the near-isogenic line of *nog1*) at the seedling stage, it was found that *NOG1* regulates the cold tolerance in rice. Using whole transcriptome analysis of near-isogenic lines, 5,954 up-regulated DEGs and 5,612 down-regulated DEGs were observed in NIL176, whereas 4,499 up-regulated DEGs and 5,850 down-regulated DEGs were found in Guichao 2 after cold treatment. The number of up-regulated genes in NIL176 was significantly higher than that in Guichao 2, among which 2,491 genes were only up-regulated in NIL176. GO annotation and KEGG enrichment analysis of DEGs showed that fatty acid metabolism, transmembrane transporter activity, integral component of membrane and plant-pathogen interaction pathways DEGs were enriched.

Materials and methods

Plant materials

The sources of NIL176 and Guichao 2 were introduced in previous studies (Huo et al., 2017). The near-isogenic line NIL176 was selected from BC₂F₂, using NIL176 as a donor and Guichao 2 as the recipient. A collection of 158 rice materials, including 84 indica (*O. sativa* L. ssp. *indica*) varieties and 74 japonica (*O. sativa* L. ssp. *japonica*) varieties, were used to verify the alleles of *NOG1* and to investigate the geographic distribution of material origins, as listed in Supplementary Table S1.

Primers

The primers used in this study for *NOG1* sequencing are listed in Supplementary Table S2.

Cold stress identification at the seedling stage

The seeds of the test materials were treated at 45°C for 48 h to break dormancy. Soak seeds with ddH₂O at room temperature for 24 h, keep moist for 24–48 h to promote germination. Seeds with uniform whiteness were selected and sown in bottomless 96-well PCR plates. Plant 3 plates of each material, hydroponically in an artificial climate chamber. The temperature of the artificial climate chamber is 28°C, with a light/darkness regime of 12/12 h and a relative humidity of 85%. The tested materials were grown to the 3-leaf stage and subjected to low temperature treatment in the artificial climate chamber. The treatment temperature was 10°C, 12 h of light, 12 h of darkness, and a relative humidity of 85%. After 10 days of treatment, growth was recovered at 28°C for 3 days and the phenotype was investigated.

RNA-sequencing and data analyses

Total RNA was extracted from the leaves of the plant with cold-treatment for 12 h using the QIAGEN plant RNA kit (Hilden, Germany). The concentration and quality of RNA were evaluated using NanoDrop 2000 UV-VIS spectrophotometer (NanoDrop Technologies, Wilmington, DE, United States). Paired-end reads were generated on a HiSeq2000 platform following the manufacturer's instructions (Illumina, United States). The analysis of RNA-sequencing (RNA-seq) data and differentially expressed genes (DEGs) were carried out as previously reported (Peng et al., 2021). Briefly, RNA-seq data were mapped on the reference genome with HISAT2 2.1.0 (Kim et al., 2019). FeatureCounts 1.6.2 was used to count the number of reads mapped on exons (Liao et al., 2014). DEGs were evaluated by edgeR 3.32.0 (Robinson et al., 2010). Genes with $p < 0.05$ and log₂ fold-changes > 1 were considered as DEGs. Further screening among the initial DEGs was performed based on fragments per kilo-base per million fragments mapped (FPKM) values. Afterward, the GO and KEGG annotations of DEGs were conducted by pannzer2 (Toronen et al., 2018) and KEGG Automatic Annotation Server (Moriya et al., 2007).

Data availability statement

The raw sequence data reported in this study have been deposited in the Genome Sequence Archive (Genomics, Proteomics & Bioinformatics 2021) in National Genomics

Data Center (Nucleic Acids Res 2022), China National Center for Bioinformation/Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA008173) that are publicly accessible at <https://ngdc.cncb.ac.cn/gsa> (Chen et al., 2021).

Author contributions

FW and WL designed and supervised this work. XH, DL, and XP analyzed the data and wrote the article. JX constructed near-isogenic lines and performed cold stress. XH, JX, YLL, and JL investigated the distribution and genotype of 158 rice varieties. XH, MZ, LK, and XZ performed RNA extraction and detection. XH, YHL, XZ, ZM, and CF participated in the revision of the manuscript. All authors reviewed and approved the final manuscript.

Funding

This work was supported by National Natural Science Foundation of China (32071992), Guangdong Basic and Applied Basic Research Foundation (2019A1515011903), Special fund for scientific innovation strategy-construction of high level Academy of Agriculture Science (R2018YJ-YB1001), The Discipline Team of Agricultural Competitive Industries in Guangdong Academy of

Agricultural Sciences (202101TD), and Guangdong Key Laboratory of New Technology in Rice Breeding (2020B1212060047).

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2022.1039677/full#supplementary-material>

References

- Ariizumi, T., Kishitani, S., Inatsugi, R., Nishida, I., Murata, N., and Toriyama, K. (2002). An increase in unsaturation of fatty acids in phosphatidylglycerol from leaves improves the rates of photosynthesis and growth at low temperatures in transgenic rice seedlings. *Plant Cell Physiol.* 43, 751–758. doi:10.1093/pcp/pcf087
- Bashir, K., Matsui, A., Rasheed, S., and Seki, M. (2019). Recent advances in the characterization of plant transcriptomes in response to drought, salinity, heat, and cold stress. *Front. Plant Sci.* 10, 658. doi:10.3389/fpls.2019.00658
- Chen, T., Chen, X., Zhang, S., Zhu, J., Tang, B., Wang, A., et al. (2021). The genome sequence archive family: toward explosive data growth and diverse data types. *Genomics Proteomics Bioinforma.* 19, 578–583. doi:10.1016/j.gpb.2021.08.001
- Fujino, K., and Matsuda, Y. (2010). Genome-wide analysis of genes targeted by qltg3-1 controlling low-temperature germinability in rice. *Plant Mol. Biol.* 72, 137–152. doi:10.1007/s11103-009-9559-x
- Fujino, K., Sekiguchi, H., Matsuda, Y., Sugimoto, K., Ono, K., and Yano, M. (2008). Molecular identification of a major quantitative trait locus, qltg3-1, controlling low-temperature germinability in rice. *Proc. Natl. Acad. Sci. U. S. A.* 105, 12623–12628. doi:10.1073/pnas.0805303105
- Hua, L., Wang, D. R., Tan, L., Fu, Y., Liu, F., Xiao, L., et al. (2015). Laba1, a domestication gene associated with long, barbed awns in wild rice. *Plant Cell* 27, 1875–1888. doi:10.1105/tpc.15.00260
- Huo, X., Wu, S., Zhu, Z., Liu, F., Fu, Y., Cai, H., et al. (2017). Nog1 increases grain production in rice. *Nat. Commun.* 8, 1497. doi:10.1038/s41467-017-01501-8
- Ito, V. C., and Lacerda, L. G. (2019). Black rice (*Oryza sativa* L.): A review of its historical aspects, chemical composition, nutritional and functional properties, and applications and processing technologies. *Food Chem.* 301, 125304. doi:10.1016/j.foodchem.2019.125304
- Jaglo, K. R., Kleff, S., Amundsen, K. L., Zhang, X., Haake, V., Zhang, J. Z., et al. (2001). Components of the Arabidopsis C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in Brassica napus and other plant species. *Plant Physiol.* 127, 910–917. doi:10.1104/pp.010548
- Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* 37, 907–915. doi:10.1038/s41587-019-0201-4
- Konishi, S., Izawa, T., Lin, S. Y., Ebana, K., Fukuta, Y., Sasaki, T., et al. (2006). An SNP caused loss of seed shattering during rice domestication. *Science* 312, 1392–1396. doi:10.1126/science.1126410
- Lesk, C., Rowhani, P., and Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature* 529, 84–87. doi:10.1038/nature16467
- Li, C., Zhou, A., and Sang, T. (2006). Rice domestication by reducing shattering. *Science* 311, 1936–1939. doi:10.1126/science.1123604
- Li, C., Liu, J., Bian, J., Jin, T., Zou, B., Liu, S., et al. (2021). Identification of cold tolerance QTLs at the bud burst stage in 211 rice landraces by GWAS. *BMC Plant Biol.* 21, 542. doi:10.1186/s12870-021-03317-7
- Liao, Y., Smyth, G. K., and Shi, W. (2014). FeatureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930. doi:10.1093/bioinformatics/btt656
- Lin, S. Z., Zhang, Z. Y., Lin, Y. Z., Zhang, Q., and Guo, H. (2004). The role of calcium and calmodulin in freezing-induced freezing resistance of Populus tomentosa cuttings. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Bao* 30, 59–68.
- Lissarre, M., Ohta, M., Sato, A., and Miura, K. (2010). Cold-responsive gene regulation during cold acclimation in plants. *Plant Signal. Behav.* 5, 948–952. doi:10.4161/psb.5.8.12135
- Liu, F., Xu, W., Song, Q., Tan, L., Liu, J., Zhu, Z., et al. (2013). Microarray-assisted fine-mapping of quantitative trait loci for cold tolerance in rice. *Mol. Plant* 6, 757–767. doi:10.1093/mp/sss161
- Ma, Y., Dai, X., Xu, Y., Luo, W., Zheng, X., Zeng, D., et al. (2015). Cold1 confers chilling tolerance in rice. *Cell* 160, 1209–1221. doi:10.1016/j.cell.2015.01.046

- Manishankar, P., and Kudla, J. (2015). Cold tolerance encoded in one snp. *Cell* 160, 1045–1046. doi:10.1016/j.cell.2015.02.037
- Mao, D., Xin, Y., Tan, Y., Hu, X., Bai, J., Liu, Z. Y., et al. (2019). Natural variation in the *han1* gene confers chilling tolerance in rice and allowed adaptation to a temperate climate. *Proc. Natl. Acad. Sci. U. S. A.* 116, 3494–3501. doi:10.1073/pnas.1819769116
- Miura, K., and Furumoto, T. (2013). Cold signaling and cold response in plants. *Int. J. Mol. Sci.* 14, 5312–5337. doi:10.3390/ijms14035312
- Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C., and Kanehisa, M. (2007). Kaas: an automatic genome annotation and pathway reconstruction server. *Nucleic Acids Res.* 35, W182–W185. doi:10.1093/nar/gkm321
- Peng, X., Tun, W., Dai, S. F., Li, J. Y., Zhang, Q. J., Yin, G. Y., et al. (2021). Genome-wide analysis of cct transcript factors to identify genes contributing to photoperiodic flowering in *oryza rufipogon*. *Front. Plant Sci.* 12, 736419. doi:10.3389/fpls.2021.736419
- Pradhan, S. K., Pandit, E., Nayak, D. K., Behera, L., and Mohapatra, T. (2019). Genes, pathways and transcription factors involved in seedling stage chilling stress tolerance in indica rice through rna-seq analysis. *BMC Plant Biol.* 19, 352. doi:10.1186/s12870-019-1922-8
- Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). Edger: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140. doi:10.1093/bioinformatics/btp616
- Saijo, Y., and Loo, E. P. (2020). Plant immunity in signal integration between biotic and abiotic stress responses. *New Phytol.* 225, 87–104. doi:10.1111/nph.15989
- Saito, K., Hayano-Saito, Y., Maruyama-Funatsuki, W., Sato, Y., and Kato, A. (2004). Physical mapping and putative candidate gene identification of a quantitative trait locus *ctb1* for cold tolerance at the booting stage of rice. *Theor. Appl. Genet.* 109, 515–522. doi:10.1007/s00122-004-1667-z
- Sakamoto, T., and Murata, N. (2002). Regulation of the desaturation of fatty acids and its role in tolerance to cold and salt stress. *Curr. Opin. Microbiol.* 5, 208–210. doi:10.1016/s1369-5274(02)00306-5
- Shi, Y., and Gong, Z. (2015). One snp in *cold1* determines cold tolerance during rice domestication. *J. Genet. Genomics.* 42, 133–134. doi:10.1016/j.jgg.2015.03.007
- Su, C. F., Wang, Y. C., Hsieh, T. H., Lu, C. A., Tseng, T. H., and Yu, S. M. (2010). A novel *mybs3*-dependent pathway confers cold tolerance in rice. *Plant Physiol.* 153, 145–158. doi:10.1104/pp.110.153015
- Sweeney, M. T., Thomson, M. J., Pfeil, B. E., and McCouch, S. (2006). Caught red-handed: *rc* encodes a basic helix-loop-helix protein conditioning red pericarp in rice. *Plant Cell* 18, 283–294. doi:10.1105/tpc.105.038430
- Tan, L., Li, X., Liu, F., Sun, X., Li, C., Zhu, Z., et al. (2008). Control of a key transition from prostrate to erect growth in rice domestication. *Nat. Genet.* 40, 1360–1364. doi:10.1038/ng.197
- Toronen, P., Medlar, A., and Holm, L. (2018). Pannzer2: a rapid functional annotation web server. *Nucleic Acids Res.* 46, W84–W88. doi:10.1093/nar/gky350
- Wai, A. H., Cho, L. H., Peng, X., Waseem, M., Lee, D. J., Lee, J. M., et al. (2021). Genome-wide identification and expression profiling of alba gene family members in response to abiotic stress in tomato (*solanum lycopersicum* l.). *BMC Plant Biol.* 21, 530. doi:10.1186/s12870-021-03310-0
- Zhang, Q., Chen, Q., Wang, S., Hong, Y., and Wang, Z. (2014). Rice and cold stress: methods for its evaluation and summary of cold tolerance-related quantitative trait loci. *Rice* 7, 24. doi:10.1186/s12284-014-0024-3
- Zhang, J., Li, X. M., Lin, H. X., and Chong, K. (2019). Crop improvement through temperature resilience. *Annu. Rev. Plant Biol.* 70, 753–780. doi:10.1146/annurev-arplant-050718-100016
- Zhao, H., Kosma, D. K., and Lu, S. (2021). Functional role of long-chain acyl-coa synthetases in plant development and stress responses. *Front. Plant Sci.* 12, 640996. doi:10.3389/fpls.2021.640996
- Zhu, B. F., Si, L., Wang, Z., Zhou, Y., Zhu, J., Shangguan, Y., et al. (2011). Genetic control of a transition from black to straw-white seed hull in rice domestication. *Plant Physiol.* 155, 1301–1311. doi:10.1104/pp.110.168500
- Zhu, Z., Tan, L., Fu, Y., Liu, F., Cai, H., Xie, D., et al. (2013). Genetic control of inflorescence architecture during rice domestication. *Nat. Commun.* 4, 2200. doi:10.1038/ncomms3200